

**NON-PERTURBATIVE RENORMALISATION OF THE
LATTICE $\Delta S = 2$ FOUR-FERMION OPERATOR****A. Donini^a, G. Martinelli^{b,*}, C.T. Sachrajda^c, M. Talevi^a and A. Vladikas^d**^a Dip. di Fisica, Università degli Studi di Roma “La Sapienza” and
INFN, Sezione di Roma, P.le A. Moro 2, 00185 Rome, Italy.^b Theory Division, CERN, 1211 Geneva 23, Switzerland.^c Dep. of Physics, University of Southampton,
Southampton SO17 1BJ, U.K.^d INFN, Sezione di Roma II and
Dip. di Fisica, Università di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, 00133 Rome, Italy.**Abstract**

We compute the renormalised four-fermion operator $O^{\Delta S=2}$ using a non-perturbative method recently introduced for determining the renormalisation constants of generic lattice composite operators. Because of the presence of the Wilson term, $O^{\Delta S=2}$ mixes with operators of different chiralities. A projection method to determine the mixing coefficients is implemented. The numerical results for the renormalisation constants have been obtained from a simulation performed using the SW-Clover quark action, on a $16^3 \times 32$ lattice, at $\beta = 6.0$. We show that the use of the constants determined non-perturbatively improves the chiral behaviour of the lattice kaon matrix element $\langle \bar{K}^0 | O^{\Delta S=2} | K^0 \rangle_{\text{latt}}$.

CERN-TH/95-23
August 1995

* On leave of absence from Dip. di Fisica, Università degli Studi “La Sapienza”, Rome, Italy.

1 Introduction

Renormalisation of lattice operators is a necessary step for obtaining physical results from numerical simulations. In this paper, we apply the general method introduced in [1] to the four-fermion operator¹

$$O^{\Delta S=2} = (\bar{s}\gamma_\mu^L d)(\bar{s}\gamma_\mu^L d), \quad (1)$$

which appears in the weak effective Hamiltonian relevant for $K^0-\bar{K}^0$ mixing

$$\mathcal{H}_{\text{eff}}^{\Delta S=2} = C(M_W/\mu)O^{\Delta S=2}(\mu), \quad (2)$$

where $\gamma_\mu^L = \frac{1}{2}\gamma_\mu(1 - \gamma_5)$, $O^{\Delta S=2}(\mu)$ is the renormalised operator, $C(M_W/\mu)$ is the corresponding Wilson coefficient and μ the renormalisation scale. The $K^0-\bar{K}^0$ matrix element of $O^{\Delta S=2}(\mu)$ defines the so-called kaon B -parameter

$$\langle \bar{K}^0 | O^{\Delta S=2}(\mu) | K^0 \rangle = \frac{8}{3} f_K^2 m_K^2 B_K(\mu). \quad (3)$$

The uncertainty in the value of this matrix element restricts the precision with which the CKM matrix elements ρ and η (in the Wolfenstein parametrisation) can be determined from experimental measurements. It is therefore of considerable importance to determine this matrix element using lattice simulations.

In the continuum, chiral symmetry implies that the kaon matrix element of $O^{\Delta S=2}$ vanishes in the chiral limit [2, 3]

$$\langle \bar{K}^0(q) | O^{\Delta S=2}(\mu) | K^0(p) \rangle = \gamma(p \cdot q) + O((p \cdot q)^2). \quad (4)$$

On the lattice however, in simulations based on Wilson's formulation of the fermion action (such as the standard Wilson action or the SW-Clover action [4]), the presence of chiral symmetry breaking terms leads to the mixing of $O^{\Delta S=2}$ with operators of different chirality [5]–[10], and the matrix element of $O^{\Delta S=2}$ is different from zero at $p \cdot q = 0$ [11]–[17]. For this reason, it is possible to define a renormalised operator with definite chiral properties only in the continuum limit, i.e. when $a \rightarrow 0$. At finite a , one can improve the chiral behaviour of the matrix element of $O^{\Delta S=2}$, by subtracting a suitable set of dimension six operators. The mixing coefficients have so far been computed only in one-loop perturbation theory [5],[8]–[10]. In this way, the systematic error in the value of the matrix element determined on the lattice is of $O(\alpha_s^2)$. In addition, as a consequence of the finiteness of the lattice spacing, there are errors of $O(a)$. Following Symanzik's proposal, one can reduce these discretization errors from $O(a)$ to $O(\alpha_s a)$ by using the tree-level “improved” SW-Clover lattice quark action [4, 18]. Using this action, the improvement has been shown to be effective for two-fermion operators, at values of β currently used in numerical simulations [19]–[22]. It remains true however, that ignorance of higher-order perturbative corrections to the mixing coefficients can distort the chiral behaviour of the operator and hence induce a large systematic error in the determination of B_K . The use of a non-perturbative approach to the determination of the renormalisation constants, should reduce this systematic effect.

¹We use the Euclidean metric throughout this paper.

In the following, we will define a renormalised operator $O^{\Delta S=2}(\mu)$, obtained by applying the non-perturbative method of ref. [1] to the computation of the mixing coefficients and of the overall renormalisation constant. In order to reduce the discretisation errors, including those induced by the mixing with higher dimensional operators, it is necessary to use an improved fermion action and operators. In the computations described below we have used the improved SW-Clover action and the “improved-improved” operators introduced in ref. [10, 20]. We monitor the effects of the non-perturbative determination of the mixing coefficients by comparing the chiral behaviour of the matrix element $\langle \bar{K}^0(q) | O^{\Delta S=2} | K^0(p) \rangle_{\text{latt}}$ computed by using the operator renormalised with standard or boosted [23] perturbation theory to the matrix element of the operator renormalised non-perturbatively. In particular, by parametrizing the matrix element near the chiral limit in the standard way [12]–[15],

$$\langle \bar{K}^0 | O^{\Delta S=2} | K^0 \rangle_{\text{latt}} = \alpha + \beta m_K^2 + \gamma(p \cdot q) + \delta m_K^4 + \epsilon m_K^2(p \cdot q) + \zeta(p \cdot q)^2 + \dots, \quad (5)$$

we investigate the differences in the values of α and β , obtained by fitting the dependence of the matrix element on the kaon masses and momenta. Since in the continuum α and β are absent, cf. eq. (4), we consider a reduction of their values as a measure of the improvement in the chiral behaviour and in the accuracy of the determination of the matrix element. Using the data of the APE collaboration [16, 17], we show that the chiral behaviour is indeed improved by using the non-perturbative results.

The paper is organized as follows. In section 2, we briefly summarize the strategy followed for computing the mixing coefficients and the overall renormalisation constant of the relevant four-fermion operator; in section 3, we illustrate the projection method used to determine the mixing coefficients. Although the method is applied specifically to the renormalisation of the operator $O^{\Delta S=2}$, it can readily be generalised to other sets of operators which mix under renormalisation. In section 4 we give some information about the perturbative evaluation of the renormalisation constants on the lattice; in section 5 we present the details of the numerical simulation and discuss our results and, finally, we present our conclusions in section 6.

2 The non-perturbative method for four-fermion operators

The renormalisation method proposed in [1] completely avoids the use of lattice perturbation theory and allows for a non-perturbative determination of the renormalisation constants of any composite operator in a renormalisation scheme which is independent of the method used to regulate the ultra-violet divergences. In particular, the renormalised operators are independent of the fact that we start from bare operators in lattice QCD. To stress this point further we will refer to the renormalisation scheme defined below for $O^{\Delta S=2}$ as the RI (Regularization Independent) scheme [24]². Non-perturbative renormalisation conditions are imposed directly on quark Green functions with off-shell external states in a fixed gauge,

² Although of course such a name could be applied equally well to many other schemes.

for example the Landau gauge. The method is expected to work in all cases where it is possible to fix the virtuality of the external states $p^2 = \mu^2$ so as to satisfy the condition $\Lambda_{\text{QCD}} \ll \mu \ll 1/a$. The condition $\mu \gg \Lambda_{\text{QCD}}$ is necessary because one has to match perturbatively the effective Hamiltonian, expressed in terms of operators renormalised at the scale μ , to the full theory. This condition is common to all approaches currently used. The condition $\mu \ll 1/a$ is a requirement common to all lattice methods and is due to the presence of $O(a)$ ($O(\alpha_s a)$) effects in the operator matrix elements. The existence of the “window” $\Lambda_{\text{QCD}} \ll \mu \ll 1/a$ depends on the value of the bare lattice coupling β at which the numerical calculations are performed. We refer the reader to ref. [1] for a more detailed discussion on this point.

In the following, in order to use a more transparent notation in the formulae, we will consider the operator

$$O_+ = \frac{1}{2}[(\bar{\psi}_1 \gamma_\mu^L \psi_2)(\bar{\psi}_3 \gamma_\mu^L \psi_4) + (2 \leftrightarrow 4)], \quad (6)$$

with four distinct quark flavours ($f = 1, 2, 3, 4$) instead of the operator $O^{\Delta S=2}$. O_+ and $O^{\Delta S=2}$ have the same renormalisation properties.

The discretization of the quark action *à la* Wilson, induces a mixing of the operator (6) with operators of a different chirality which, in the language of refs. [24]–[27], correspond to the so-called “effervescent” (“evanescent”) operators. The mixing, being a consequence of the regularization procedure, is not limited to the lattice case, but is present also in continuum regularizations. The effervescent operators must be subtracted from the bare one by a suitable renormalisation procedure³.

In the lattice case, CPS symmetry fixes the basis of operators that may appear in perturbation theory [28]:

$$O_+^{SP} = -\frac{1}{16N_c}[(\bar{\psi}_1 \psi_2)(\bar{\psi}_3 \psi_4) - (\bar{\psi}_1 \gamma_5 \psi_2)(\bar{\psi}_3 \gamma_5 \psi_4) + (2 \leftrightarrow 4)], \quad (7)$$

$$O_+^{VA} = -\frac{(N_c^2 + N_c - 1)}{32N_c}[(\bar{\psi}_1 \gamma_\mu \psi_2)(\bar{\psi}_3 \gamma_\mu \psi_4) - (\bar{\psi}_1 \gamma_\mu \gamma_5 \psi_2)(\bar{\psi}_3 \gamma_\mu \gamma_5 \psi_4) + (2 \leftrightarrow 4)], \quad (8)$$

$$O_+^{SPT} = \frac{(N_c - 1)}{16N_c}[(\bar{\psi}_1 \psi_2)(\bar{\psi}_3 \psi_4) + (\bar{\psi}_1 \gamma_5 \psi_2)(\bar{\psi}_3 \gamma_5 \psi_4) + (\bar{\psi}_1 \sigma_{\mu\nu} \psi_2)(\bar{\psi}_3 \sigma_{\mu\nu} \psi_4) + (2 \leftrightarrow 4)], \quad (9)$$

where $N_c = 3$ denotes the number of colours.

We renormalise the operator O_+ by introducing the subtracted operator O_+^s ,

$$O_+(\mu) = Z_+ O_+^s = Z_+(O_+ + Z_1 O_+^{SP} + Z_2 O_+^{VA} + Z_3 O_+^{SPT}), \quad (10)$$

where $Z_{+,1,2,3} = Z_{+,1,2,3}(\mu a, g_0^2(a))$ and the bare lattice coupling is given by the relation $\beta = 6/g_0^2(a)$. The mixing constants Z_i , $i = 1, \dots, 3$, are determined by means of projection

³ Using dimensional regularization, the one-loop mixing with the “effervescent” operators is cancelled by the minimal subtraction of the pole in $1/\epsilon$.

operators that will be defined in section 3. Their values are fixed by the requirement that, up to terms of $O(\alpha_s a)$, O_+^s renormalises multiplicatively. O_+^s is logarithmically divergent as $a \rightarrow 0$, and this divergence is removed by imposing a renormalisation condition on O_+^s which defines the overall renormalisation constant $Z_+(\mu a, g_0^2(a))$,

$$Z_+(\mu a, g_0^2(a)) Z_\psi^{-2}(\mu a, g_0^2(a)) \Gamma_+^s(pa)|_{p^2=\mu^2} = 1, \quad (11)$$

where $\Gamma_+^s(pa)$ is obtained by projecting a suitable amputated Green function of the operator O_+^s on the Dirac structure $\gamma_\mu^L \otimes \gamma_\mu^L$ (see eq. (23) in section 3 and refs. [1, 24]). Z_ψ is the quark field renormalisation constant to be defined below (eq. (33) of section 3 and ref. [1]). In eq. (11), $p^2 = \mu^2$ denotes the momentum of the external quark states. We have chosen equal momenta for all four external quark legs, because this is the simplest choice which regulates the infrared divergences [24].

The renormalised operator in eq. (10), calculated in the RI scheme, depends both on the gauge and on the external states. The Wilson coefficient must also be calculated in the same gauge and with the same external states in order to obtain the physical operators which are independent of both⁴. The next-to-leading order calculation of the Wilson coefficient relevant for the operator (6), in the Landau gauge and with equal external momenta, can be found in ref. [24].

3 Determination of the mixing constants

In this section, we define the four-point amputated Green functions and introduce the projectors that have been used to determine the mixing constants.

Since the non-perturbative renormalisation conditions are imposed on quark states, the Green functions of a four-fermion operator will depend on four coordinates. Denoting by x_1, x_3 and x_2, x_4 the coordinates of the outgoing and incoming quarks, the Green functions corresponding to the insertion of the operators (6)–(9) can be written as linear combinations of Green functions of the form

$$G_{\Gamma^a}(x_1, x_2, x_3, x_4) = \langle \psi_1(x_1) \bar{\psi}_2(x_2) O_{\Gamma^a}(0) \psi_3(x_3) \bar{\psi}_4(x_4) \rangle, \quad (12)$$

where $\langle \dots \rangle$ denotes the vacuum expectation value, i.e. the average over the gauge-field configurations. The generic four-fermion operator O_{Γ^a} is given by

$$O_{\Gamma^a}(0) = C_{\Gamma^a} \left[\bar{\psi}_1(0) \Gamma^a \psi_2(0) \bar{\psi}_3(0) \Gamma^a \psi_4(0) + \bar{\psi}_1(0) \Gamma^a \psi_4(0) \bar{\psi}_3(0) \Gamma^a \psi_2(0) \right], \quad (13)$$

where Γ^a denotes a Dirac matrix, and C_{Γ^a} is a constant associated with Γ^a . The index a can be either single-valued (if $\Gamma^a = \mathbf{1}$ or γ_5) or be summed over a range of values (if $\Gamma^a = \gamma_\mu$, $\gamma_\mu \gamma_5$ or $\sigma_{\mu\nu}$ a sum over repeated Lorentz indices is implied).

The Fourier transform of the non-amputated Green function (12), at equal external momenta p , has the form

$$G_{\Gamma^a}(p)_{\alpha\beta\gamma\delta}^{ABCD} = C_{\Gamma^a} \left[\langle \Gamma^a(p)_{\alpha\beta}^{AB} \otimes \Gamma^a(p)_{\gamma\delta}^{CD} \rangle - \langle \Gamma^a(p)_{\alpha\delta}^{AD} \otimes \Gamma^a(p)_{\gamma\beta}^{CB} \rangle \right], \quad (14)$$

⁴This is true up to higher order continuum perturbative corrections and lattice systematic errors.

where

$$\Gamma^a(p)_{\chi\psi}^{XY} = S(p|0)_{\chi\rho}^{XR} \Gamma_{\rho\sigma}^a (\gamma_5 S(p|0)^\dagger \gamma_5)_{\sigma\psi}^{RY}. \quad (15)$$

In eqs. (14) and (15) the upper-case Roman superscripts denote colour labels and the lower case Greek subscripts denote spinor labels. $S(p|0)$ is defined by

$$S(p|0) = \int d^4x S(x|0) e^{-ip \cdot x}, \quad (16)$$

where $S(x|0)$ is the quark propagator computed on a single gauge-field configuration (cf. section 4 of [1]), and is therefore not translationally invariant. It satisfies the relation

$$S(x|0) = \gamma_5 S^\dagger(0|x) \gamma_5. \quad (17)$$

The amputated Green function can be obtained from eq. (14)

$$\Lambda_{\Gamma^a}(p)_{\rho\sigma\rho'\sigma'}^{RSR'S'} = S^{-1}(p)_{\rho\alpha}^{RA} S^{-1}(p)_{\rho'\gamma}^{R'C} G_{\Gamma^a}(p)_{\alpha\beta\gamma\delta}^{ABCD} S^{-1}(p)_{\beta\sigma}^{BS} S^{-1}(p)_{\delta\sigma'}^{DS'}, \quad (18)$$

where $S(p)$ is the Fourier transform of the translationally-invariant quark propagator, i.e. the Fourier transform of $S(x|0)$, averaged over the gauge-field configurations.

As mentioned above, the renormalisation procedure necessary to determine the mixing constants consists in defining suitable projectors on the amputated Green functions of the operators (6)–(9). To this end let us introduce a more convenient notation. Let us denote by O_i , $i = 0, \dots, 3$, respectively, the operators O_+ , O_+^{SP} , O_+^{VA} , O_+^{SPT} . Then, the projectors $\hat{P}_i$, $i = 0, \dots, 3$, are defined by the condition

$$\text{Tr } \hat{P}_i \Lambda_j^{(0)} = \delta_{ij}, \quad i, j = 0, \dots, 3, \quad (19)$$

where $\Lambda_i^{(0)}$, $i = 0, \dots, 3$, are the amputated Green functions, at tree level, of the operators O_i , and the trace is understood over colour and spin (as defined below). The renormalisation scheme depends on the precise definition of the projection operators and we now define our procedure in detail.

For each Dirac Matrix Γ^b , we define the projector $\hat{P}_{\Gamma^b}$ by ⁵

$$\text{Tr } \hat{P}_{\Gamma^b} \Lambda_{\Gamma^a}(p) = (\Gamma_{\sigma\rho}^b \otimes \Gamma_{\sigma'\rho'}^b) \Lambda_{\Gamma^a}(p)_{\rho\sigma\rho'\sigma'}^{RRR'R'}, \quad (20)$$

where the index b is either fixed or corresponds to a sum over repeated indices. In the free theory, the amputated Green function reduces to

$$\Lambda_{\Gamma^a}^{(0)}(p)_{\rho\sigma\rho'\sigma'}^{RSR'S'} = C_{\Gamma^a} [\delta^{RS} \delta^{R'S'} (\Gamma_{\rho\sigma}^a \otimes \Gamma_{\rho'\sigma'}^a) - \delta^{RS'} \delta^{R'S} (\Gamma_{\rho\sigma'}^a \otimes \Gamma_{\rho'\sigma}^a)]. \quad (21)$$

and the result of the projection defined in (20) is:

$$\text{Tr } \hat{P}_{\Gamma^b} \Lambda_{\Gamma^a}^{(0)}(p) = C_{\Gamma^a} [N_c^2 (\text{Tr } \Gamma^a \Gamma^b) (\text{Tr } \Gamma^a \Gamma^b) - N_c (\text{Tr } \Gamma^a \Gamma^b \Gamma^a \Gamma^b)]. \quad (22)$$

⁵It is only the traces (20) which are required for the determination of the subtraction constants.

The projectors corresponding to O_+ and to the operators O_i defined in eqs.(7)–(9) are as follows:

$$\hat{\mathbf{P}}_0 = \frac{1}{8N_c(N_c + 1)}\hat{\mathbf{P}}_{\gamma_\mu^R}, \quad (23)$$

$$\begin{aligned} \hat{\mathbf{P}}_1 &= \frac{N_c}{2(1 - N_c^2)}(\hat{\mathbf{P}}_{\mathbf{I}} - \hat{\mathbf{P}}_{\gamma_5}) \\ &+ \frac{1}{4(1 - N_c^2)}(\hat{\mathbf{P}}_{\gamma_\mu} - \hat{\mathbf{P}}_{\gamma_\mu\gamma_5}), \end{aligned} \quad (24)$$

$$\begin{aligned} \hat{\mathbf{P}}_2 &= \frac{1}{2(1 - N_c^2)(N_c^2 + N_c - 1)}(\hat{\mathbf{P}}_{\mathbf{I}} - \hat{\mathbf{P}}_{\gamma_5}) \\ &+ \frac{N_c}{4(1 - N_c^2)(N_c^2 + N_c - 1)}(\hat{\mathbf{P}}_{\gamma_\mu} - \hat{\mathbf{P}}_{\gamma_\mu\gamma_5}) \end{aligned} \quad (25)$$

$$\hat{\mathbf{P}}_3 = \frac{1}{8(N_c^2 - 1)}(\hat{\mathbf{P}}_{\mathbf{I}} + \hat{\mathbf{P}}_{\gamma_5} + \hat{\mathbf{P}}_{\sigma_{\mu\nu}}), \quad (26)$$

where $\gamma_\mu^R = \frac{1}{2}\gamma_\mu(1 - \gamma_5)$. Note that the projectors $\hat{\mathbf{P}}_0$ and $\hat{\mathbf{P}}_3$, eqs. (23) and (26), have the same Dirac structure as the operators O_0 and O_3 , eqs. (6) and (9). This is due to the Fierz rearrangement properties of these operators.

It is possible to determine the mixing coefficients Z_i by using the projectors (23)–(26) defined in the free field case. Let us introduce the matrix D defined by

$$\Lambda_i = \sum_{j=0}^3 D_{ij}\Lambda_j^{(0)}, \quad (27)$$

where the elements D_{ik} are determined non-perturbatively by the projections

$$D_{ik} = \text{Tr } \hat{\mathbf{P}}_k \Lambda_i, \quad i, k = 0, \dots, 3. \quad (28)$$

The mixing constants Z_i are then fixed by the condition that the subtracted operator O_+^s is proportional to the bare free operator

$$\text{Tr } \hat{\mathbf{P}}_k \Lambda_+^s = \left(D_{0k} + \sum_{i=1}^3 Z_i D_{ik} \right) = 0, \quad k = 1, 2, 3. \quad (29)$$

Equation (29) yields three conditions corresponding to a linear non-homogeneous system in the three unknowns Z_i . Defining the reduced 3×3 matrix $\tilde{D}$ as

$$\tilde{D}_{ik} = D_{ki}, \quad i, k = 1, 2, 3, \quad (30)$$

the solutions of this linear system are given by

$$Z_i = - \sum_{k=1}^3 (\tilde{D})_{ik}^{-1} D_{0k}, \quad i = 1, 2, 3. \quad (31)$$

The overall renormalisation constant Z_+ is then determined by the condition (11), using

$$\Gamma_+^s(pa) = \text{Tr } \hat{P}_0 \Lambda_+^s = \left(D_{00} + \sum_{i=1}^3 Z_i D_{i0} \right). \quad (32)$$

In eq. (11), the renormalisation constant Z_ψ is defined by the relation

$$Z_\psi(\mu a, g_0^2(a)) = \frac{1}{48} \text{Tr} \left(\Lambda_{V_\mu^L} \gamma_\mu \right) \Big|_{p^2=\mu^2} \times Z_{V^L}, \quad (33)$$

where $V^L = \bar{\psi} \gamma_\mu \psi$ is the local vector current, and Z_{V^L} its renormalisation constant which can be determined with high accuracy, by using the vector current Ward identities [21, 22, 29]. $\Lambda_{V_\mu^L}$ is defined as

$$\Lambda_{V_\mu^L}(p) = S(p)^{-1} G_{V_\mu^L}(p) S(p)^{-1}, \quad (34)$$

where $G_{V_\mu^L}(p)$ is the non-amputated two-point Green function of the local vector current, $G_{V_\mu^L}(p) = \langle \Gamma_{V_\mu^L}^L(p) \rangle$, cf. eq. (15). There are various equivalent ways to define Z_ψ , but (33) is the most natural from a non-perturbative point of view. For a more thorough discussion on the determination of Z_ψ , we refer the reader to sections 2 and 4 of ref. [1].

4 Lattice perturbation theory

We have also calculated the renormalisation constants Z_+, Z_1, Z_2 and Z_3 in one-loop perturbation theory, in order to be able to compare the results with those obtained non-perturbatively. Since the non-perturbative renormalisation condition depends on the gauge and on the external states, the perturbative calculation must be done in the Landau gauge and at equal external momenta. This calculation is an extension of those of refs. [9, 10].

Starting from a bare lattice operator $O(a)$, the one-loop vertex function $\Gamma_O^\lambda(pa)$ is obtained by tracing the amputated Green function (but with wave function effects included) with a suitable projector. The generic expression of $\Gamma_O^\lambda(pa)$, calculated between states of momentum p and in a fixed gauge λ , is

$$\Gamma_O^\lambda(pa) = \left[1 + \frac{\alpha_s}{4\pi} \left(\gamma_O \log(1/p^2 a^2) + r_O^{\text{latt}}(\lambda, p) \right) \right]. \quad (35)$$

γ_O is the anomalous dimension, which at one-loop order is independent of the gauge, the external states and the regularization⁶. The finite coefficient $r_O^{\text{latt}}(\lambda, p)$, on the other hand, does depend on the gauge, the regularization and the external states. The momentum label p appearing as argument of $r_O^{\text{latt}}(\lambda, p)$ indicates that the result is a dimensionless function of the external states.

⁶ λ denotes a generic covariant gauge: $\lambda = 0$ corresponds to the Landau gauge, $\lambda = 1$ correspond to the Feynman gauge.

In the continuum, in any renormalisation scheme based on dimensional regularization (DR=NDR, HV or DRED), the vertex function between states of momentum p and in a generic gauge, is

$$\Gamma_O^\lambda(p/\mu) = \left[1 + \frac{\alpha_s}{4\pi} \left(\gamma_O \log(\mu^2/p^2) + r_O^{\text{DR}}(\lambda, p) \right) \right], \quad (36)$$

where μ is the DR renormalisation scale. Thus, the one-loop relation between the operators in the continuum and on the lattice is

$$O(\mu) = \left[1 + \frac{\alpha_s}{4\pi} \left(\gamma_O \log(\mu^2 a^2) + \Delta^{\text{DR-latt}} \right) \right] O(a), \quad (37)$$

where

$$\Delta_O^{\text{DR-latt}} = r_O^{\text{DR}}(\lambda, p) - r_O^{\text{latt}}(\lambda, p) \quad (38)$$

is independent of both λ and p . From $\Delta_O^{\text{DR-latt}}$, we can calculate the lattice constant $r_O^{\text{latt}}(\lambda, p)$, in any gauge and at any external momenta, from the corresponding constant in the continuum, $r_O^{\text{DR}}(\lambda, p)$. In order to compare the perturbative result with the non-perturbative determination, we need $r_O^{\text{latt}}(\lambda = 0, p)$ in the Landau gauge and with non-zero but equal external momenta. From (38), we immediately obtain

$$r_O^{\text{latt}}(\lambda = 0, p) = r_O^{\text{DR}}(\lambda = 0, p) - \Delta_O^{\text{DR-latt}}. \quad (39)$$

Since r_O^{latt} must be independent of the continuum regularization used in the intermediate steps, a check of the correctness of the calculation is given by

$$r_O^{\text{DRED}}(\lambda = 0, p) - \Delta_O^{\text{DRED-latt}} = r_O^{\text{NDR}}(\lambda = 0, p) - \Delta_O^{\text{NDR-latt}}, \quad (40)$$

which is equivalent to

$$r_O^{\text{DRED}}(\lambda = 0, p) - r_O^{\text{NDR}}(\lambda = 0, p) = r_O^{\text{DRED}}(\lambda = 1, p') - r_O^{\text{NDR}}(\lambda = 1, p'). \quad (41)$$

The one-loop contribution to the renormalisation constant Z_+ and to the mixing coefficients Z_i 's have been calculated in [9, 10], by comparing the lattice and the DRED scheme. In the notation of these authors

$$\begin{aligned} Z_+ &= 1 + \frac{\alpha_s}{4\pi} F_+, & F_+ &= \Delta_{O_+}^{\text{DRED-latt}} = -10.9, \\ Z_1 &= Z_2 = Z_3 = \frac{\alpha_s}{4\pi} F^*, & F^* &= 19.4. \end{aligned} \quad (42)$$

In the DRED scheme, for a generic gauge λ and by taking the momenta of the external legs to be equal, we find

$$r_{O_+}^{\text{DRED}}(\lambda, p) = \lambda \left(-7/3 + 8/3 \log(2) \right) - 5/3 + 8 \log(2), \quad (43)$$

whilst the mixing with the “effervescent” operators is cancelled by the minimal subtraction procedure. Thus, the perturbative expressions of the lattice renormalisation constants in the RI scheme are given by⁷

$$\begin{aligned} Z_+^{\text{RI}} &= 1 - \frac{\alpha_s}{4\pi} \left(-\gamma_{O_+} \log(\mu^2 a^2) + r_{O_+}^{\text{DRED}}(\lambda = 0, p) - F_+ \right), \\ Z_i^{\text{RI}} &= \frac{\alpha_s}{4\pi} F^*, \end{aligned} \quad (44)$$

with $\gamma_{O_+} = -2$ and $r_{O_+}^{\text{DRED}}(\lambda, p)$ given in eq. (43).

We have not evaluated the renormalisation constants using Discrete Perturbation Theory (DPT), i.e. by summing only over the discrete values of momenta allowed on our finite lattice (this was done for the two-quark operators in ref. [1]). We have only evaluated the constants using standard lattice perturbation theory, in which finite lattice size effects are neglected.

In order to estimate the values of the renormalisation constants we have used the following “boosted” coupling constant α_s^V [23]

$$\alpha_s^V = \frac{1}{\langle \frac{1}{3} \text{Tr } U_P \rangle} \alpha_s^{\text{latt}} \simeq 1.68 \alpha_s^{\text{latt}} \quad (\text{at } \beta = 6.0). \quad (45)$$

as our expansion parameter and refer to the result as corresponding to Boosted Perturbation Theory (BPT). We also present the values obtained using the bare coupling α_s^{latt} , and refer to these results as coming from standard perturbation theory (SPT).

5 Numerical results

In this section, we give the numerical results of our calculation. The simulation has been performed by generating 36 independent gluon-field configurations, on a $16^3 \times 32$ lattice, at $\beta = 6.0$. The errors have been obtained with the jackknife method, by decimating three configurations at a time. The SW-Clover quark propagators have been computed at a single value of the quark mass ($m_q a \simeq 0.07$), corresponding to the hopping parameter $\kappa = 0.1425$. The quark Green functions have been computed in the lattice Landau gauge, defined by minimizing the functional

$$\text{Tr} \left[\sum_{\mu=1}^4 (U_\mu(x) + U_\mu^\dagger(x)) \right]. \quad (46)$$

Possible effects from Gribov copies have not been studied. For more details, see ref. [30].

In fig. 1, the renormalisation constants, obtained by using the prescription described in sections 2 and 3, are given as a function of the renormalisation scale $\mu^2 a^2$. We hope to find an interval of values of $\mu^2 a^2$, large enough to avoid significant non-perturbative effects and small enough to avoid large discretization errors. In ref. [1], the existence of such

⁷ The scale μ in this formula denotes the renormalization scale at which $\Gamma_O^\lambda(pa)$ is renormalized, i.e. $Z_+ \Gamma_O^\lambda(pa)|_{p^2=\mu^2} = 1$.




Figure 1: *Non-perturbative renormalisation constants of the operator O_+ as a function of $\mu^2 a^2$: (a) the overall renormalisation constant Z_+ ; (b)–(d) the mixing coefficients Z_i , $i = 1, 2, 3$. We also report the perturbative evaluation: the dashed curve is from BPT, while the solid curve is from SPT.*

$\mu^2 a^2$	Z_+	Z_1	Z_2	Z_3
0.46	0.91 ± 0.05	0.08 ± 0.14	0.34 ± 0.03	0.34 ± 0.07
0.66	0.84 ± 0.05	0.07 ± 0.09	0.33 ± 0.03	0.34 ± 0.06
0.81	0.83 ± 0.04	0.11 ± 0.07	0.31 ± 0.03	0.28 ± 0.05
0.96	0.84 ± 0.03	0.14 ± 0.07	0.30 ± 0.02	0.24 ± 0.04
1.27	0.80 ± 0.04	0.17 ± 0.05	0.29 ± 0.02	0.21 ± 0.03
1.54	0.82 ± 0.02	0.19 ± 0.04	0.27 ± 0.02	0.16 ± 0.03
1.89	0.83 ± 0.03	0.22 ± 0.05	0.30 ± 0.02	0.18 ± 0.03
2.47	0.85 ± 0.02	0.22 ± 0.06	0.33 ± 0.02	0.23 ± 0.03
SPT	0.91	0.12	0.12	0.12
BPT	0.84	0.21	0.21	0.21

Table 1: Values of Z_+ and Z_i ($i = 1, 2, 3$) for several renormalisation scales $\mu^2 a^2$. We also give the results obtained at $\mu^2 a^2 = 1$, by using “standard” perturbation theory (SPT) and “boosted” perturbation theory (BPT), using an effective coupling $\alpha_s^V = 1.68 \alpha_s^{\text{latt}}$.

a “window” in $\mu^2 a^2$ was investigated by comparing the renormalisation constants of two-quark operators computed in perturbation theory with the corresponding non-perturbative determinations on quark states, and with the results obtained by using the Ward identity method [21, 22, 29]. In most of the cases a range of acceptable values was found in the interval $0.8\text{--}0.9 \leq \mu^2 a^2 \leq 1.5\text{--}2.0$. At smaller values of $\mu^2 a^2$, in particular in the case of the axial current and of the pseudoscalar density (probably because of the presence of a pseudo-Goldstone boson contribution), the non-perturbative corrections were found to be large. For this reason, it is difficult to determine the renormalisation constant of the axial current in this way. Only at values of $\mu^2 a^2$ larger than 1.5–2.0, a surprisingly large value in our opinion, did discretization errors become clearly visible. They were signalled by the fact that the renormalisation constants computed at the same values of $\mu^2 a^2$, but with inequivalent components of the momentum p (e.g. $p \equiv 2\pi/16a(4, 4, 0, 2)$ and $p \equiv 2\pi/16a(0, 0, 0, 6)$) were found to be different [1]. This was interpreted as a signal of the breaking of the Lorentz symmetry due to lattice artefacts, see also ref. [31].

For the four-fermion operators considered in this paper, we do not have the possibility of checking the results for the renormalisation constants by the use of Ward identities, but it appears that a similar situation may also occur in this case. In the region of momenta $\mu^2 a^2 \geq 0.96$, the renormalisation constants are determined with a relatively small error (the worse case being the error of Z_1 which is about 50% at $\mu^2 a^2 = 0.96$) and the dependence on the scale is relatively weak, as can be seen from table 1 and fig. 1. We notice that the values of Z_+ and Z_2 have small statistical errors even at scales smaller than 0.96 ($\sim 10\%$ in the worst case), that Z_3 has relative errors in the range of about 15–20%, and that Z_1 has the largest relative error at all the scales considered in table 1. As for the scale dependence, Z_2 is quite stable as a function of $\mu^2 a^2$, while both Z_1 , which suffers from the largest statistical uncertainty, and Z_3 do not exhibit a very clear plateau, as can be seen in fig. 1. In particular the value of Z_1 seems to increase with the scale. With the present statistics, we cannot determine whether the variation of Z_1 and Z_3 with $\mu^2 a^2$ is real or due to

$\mu^2 a^2$	α	β	γ
0.46	0.022(16)	0.23(19)	0.78(13)
0.66	0.013(15)	0.20(17)	0.70(12)
0.81	0.012(13)	0.21(17)	0.69(12)
0.96	0.017(13)	0.21(17)	0.70(12)
1.27	0.015(13)	0.21(16)	0.66(11)
1.54	0.018(13)	0.22(16)	0.67(12)
1.89	0.023(13)	0.22(16)	0.69(12)
2.47	0.022(14)	0.23(17)	0.72(12)
SPT	-0.067(12)	0.17(15)	0.62(11)
BPT	-0.054(12)	0.17(15)	0.62(11)

Table 2: Values of the coefficients α , β and γ obtained from a linear fit of $\langle O_+ \rangle = \langle \bar{K}^0 | O^{\Delta S=2} | K^0 \rangle_{\text{latt}} / \langle P_5 \rangle^2$. The results refer to operators renormalised at different scales $\mu^2 a^2$. Values of the same parameters in standard perturbation theory and in boosted perturbation theory are also reported.

statistical fluctuations. Fortunately, as we will see below, the largest correction to the chiral behaviour comes from the operator O_2 , corresponding to Z_2 , which is very well determined. Hence the chiral behaviour of the operator O_+ is stable with respect to the uncertainties above.

In order to investigate the effects of the non-perturbative corrections, we have combined our results with the computation of the lattice matrix elements of the four-fermion operators O_0 – O_3 (6)–(9) performed in ref. [17], where a more detailed discussion of the numerical aspects can be found. Here we limit ourselves to a qualitative discussion of the results.

In fig. 2, we show the chiral behaviour of $\langle O_+ \rangle = \langle \bar{K}^0 | O^{\Delta S=2} | K^0 \rangle_{\text{latt}} / \langle P_5 \rangle^2$ with the meson at rest as a function of $X = 8/3 f_K^2 M_K^2 / \langle P_5 \rangle^2$. $\langle P_5 \rangle^2 = |\langle 0 | \bar{s} \gamma_5 d | K^0 \rangle|^2$ is the squared matrix element of the pseudoscalar density between the meson and the vacuum. The variables $\langle O_+ \rangle$ and X are particularly convenient since they can be obtained from suitable two- and three-point correlation functions without any fitting procedure [13].

An analysis of the different contributions to the matrix element of the renormalised operator shows that the largest correction comes from the operator O_2 , eq. (8), whose constant Z_2 is well determined. This contribution is much larger than that coming from O_1 (and larger than that from O_3)⁸. Thus, the uncertainty in Z_1 (and partly Z_3) has no significant consequences for the value of the kaon matrix element (3). Indeed, as can be seen from table 2, in passing from $\mu^2 a^2 = 0.66$ to $\mu^2 a^2 = 2.47$, the central value of Z_1 increases by a factor of 3, but there are no large variations in the values of α , β and γ , eq. (5), i.e. the “physical” results depend rather weakly on the scale. The use of the non-perturbative renormalisation constants leaves the values of β and γ almost unchanged compared to those obtained by using the constants computed in one-loop perturbation theory: $\beta \sim 0.2$, and within the errors is compatible with zero in both cases, and γ is about 15% larger in the non-perturbative

⁸Notice that O_1 has the smallest colour factor.




Figure 2: *Chiral behaviour of $\langle O_+ \rangle$ as a function of X (see text). We give the matrix elements of the operator renormalised in standard perturbation theory ($\diamond$), boosted perturbation theory ($\square$) and non-perturbatively ($\circ$).*

case. In contrast, α changes sign and its absolute value is reduced by about a factor of three in the non-perturbative case, and becomes compatible with zero. This happens for any choice of $\mu^2 a^2$ between 0.46 and 2.47, cf. tab. 1. The variation of the Z_i 's in the interval of $\mu^2 a^2$ considered in tables 1 and 2 is representative of the variation allowed by the statistical errors. Since the Z_i 's and the matrix elements of the four-fermion operators have been computed on different sets of configurations, this is the best test of the stability of the results which can be done at present. A more definite conclusion will be reached by computing the renormalisation constants and the matrix elements on the same set of configurations.

Since α should vanish in the continuum limit, we conclude that the use of the non-perturbative renormalisation constants improves the chiral behaviour for a large range of values of the renormalisation scale. This is also illustrated in fig. 2 [17], obtained for $\mu^2 a^2 = 0.96$. More details will be given in ref. [17].

6 Conclusions

We have applied the non-perturbative renormalisation method proposed in ref. [1] to the $\Delta S = 2$ operator given in eq. (1). Since, on the lattice, this operator mixes with other dimension-six operators of different chirality, we have illustrated a projection method for the determination of the mixing coefficients. The overall renormalisation constant of the subtracted operator has then been obtained as in the case of any other multiplicatively renormalisable operator.

In this exploratory study we have computed the subtraction constants with limited statistical precision (36 configurations on a $16^3 \times 32$ lattice at $\beta = 6.0$ using the improved SW-Clover action). The results in fig.1 and table 1 are very encouraging, and motivate us to repeat the calculation with larger statistics and at different values of β ($\beta = 6.2$ and 6.4), and to extend it to the operators relevant for $\Delta I = 1/2$ transitions and to the penguin operators which control CP-violation in kaon systems. Even with our limited statistical precision, our results indicate that the chiral behaviour of the $K^0\text{--}\bar{K}^0$ matrix element of $O^{\Delta S=2}(\mu)$ is improved significantly by the use of the subtraction constants which were determined non-perturbatively. This supports our view that, by combining the improvement of the action *à la* Symanzik, which reduces $O(a)$ effects, with the non-perturbative method of ref. [1], which reduces higher-order effects in the mixing coefficients, it is possible to achieve an accurate determination of the physical weak amplitudes using Wilson-like fermions.

Acknowledgements

We warmly thank the members of the APE collaboration for the use of their results before publication. M.T. thanks the Physics Department of Southampton University for their kind hospitality during the completion of this work. We acknowledge the partial support by the EC contract CHRX-CT92-0051 and by M.U.R.S.T., Italy. C.T.S. acknowledges the Particle Physics and Astronomy Research Council for its support through the award of a Senior Fellowship. We also acknowledge the Computer Centre of CINECA (Bologna, Italy), where these calculations were performed, and thank their staff for their precious help.

References

- [1] G. Martinelli, C. Pittori, C.T. Sachrajda, M. Testa and A. Vladikas, Nucl. Phys. B445 (1995) 81.
- [2] N. Cabibbo, Phys. Rev. Lett. 12 (1964) 62.
- [3] M. Gell-Mann, Phys. Rev. Lett. 12 (1964) 155.
- [4] B. Sheikholeslami and R. Wohlert, Nucl. Phys. B259 (1985) 572.

- [5] G. Martinelli, Phys. Lett. B141 (1984) 395.
- [6] M. Boicichio, L. Maiani, G. Martinelli, G.C. Rossi and M. Testa, Nucl. Phys. B262 (1985) 331.
- [7] L. Maiani, G. Martinelli, G.C. Rossi and M. Testa, Phys. Lett. B176 (1986) 445; Nucl. Phys. B289 (1987) 505.
- [8] C. Bernard, T. Draper and A. Soni, Phys. Rev. D36 (1987) 3224.
- [9] R. Frezzotti, E. Gabrielli, C. Pittori and G.C. Rossi, Nucl. Phys. B373 (1991) 781.
- [10] A. Borrelli, R. Frezzotti, E. Gabrielli and C. Pittori, Nucl. Phys. B409 (1993) 382.
- [11] C. Bernard, T. Draper, G. Hockney, A.M. Rushton and A. Soni, Phys. Rev. Lett. 55 (1985) 2770.
- [12] APE collaboration, presented by G. Martinelli, Nucl. Phys. B (Proc. Suppl.) 17 (1990) 523.
- [13] M.B. Gavela *et al.*, Nucl. Phys. B306 (1988) 677.
- [14] C. Bernard and A. Soni, Nucl. Phys. B (Proc. Suppl.) 17 (1990) 495; Nucl. Phys. B (Proc. Suppl.) 20 (1991) 410.
- [15] R. Gupta, D. Daniel, G.W. Kilcup, A. Patel and S. Sharpe, Phys. Rev. D46 (1993) 5113.
- [16] APE Collaboration, M. Crisafulli *et al.*, presented by A. Donini, Nucl. Phys. B (Proc. Suppl.) 42 (1995) 397.
T. Bhattacharya and R. Gupta, Nucl. Phys. B (Proc. Suppl.) 42 (1995) 935.
- [17] APE Collaboration, M. Crisafulli *et al.*, ROME Prep. 1105/95, in preparation.
- [18] G. Heatlie, G. Martinelli, C. Pittori, G.C. Rossi and C.T. Sachrajda, Nucl. Phys. B352 (1991) 266.
- [19] G. Martinelli, C.T. Sachrajda and A. Vladikas, Nucl. Phys. B358 (1991) 212.
- [20] G. Martinelli, C.T. Sachrajda, G. Salina and A. Vladikas, Nucl. Phys. B378 (1992) 591.
- [21] G. Martinelli, S. Petrarca, C.T. Sachrajda and A. Vladikas, Phys. Lett. B311 (1993) 241.
- [22] UKQCD Collaboration, D.S. Henty, R.D. Kenway, B.J. Pendleton and J.I. Skullerud, Edinburgh prep. 94/545, hep-lat 9412088.
- [23] G.P. Lepage and P.B. Mackenzie, Nucl. Phys. B (Proc. Suppl.) 20 (1991) 173; Phys. Rev. D48 (1993) 2250.

- [24] M. Ciuchini, E. Franco, G. Martinelli and L. Reina, Rome prep. 94/1024, CERN-TH 7514/94, hep-ph 9501265, to appear in Z. Phys. C.
- [25] G. Altarelli, G. Curci, G. Martinelli and S. Petrarca, Nucl. Phys. B187 (1981) 461.
- [26] A.J. Buras, M. Jamin, M.E. Lautenbacher and P.H. Weisz, Nucl. Phys. B370 (1992) 69; Addendum Nucl Phys. B375 (1992) 501.
- [27] M. Ciuchini, E. Franco, G. Martinelli and L. Reina, Nucl. Phys. B415 (1994) 403.
- [28] C. Bernard, T. Draper, G. Hockney and A. Soni, Nucl. Phys. B (Proc. Suppl.) 4 (1988) 483.
- [29] L. Maiani and G. Martinelli, Phys. Lett. B178 (1986) 265.
- [30] M.L. Paciello, S. Petrarca, B. Taglienti and A. Vladikas, Phys. Lett. B341 (1994) 187.
- [31] P. Marenzoni, G. Martinelli and N. Stella, Rome prep. 94/1042- SHEP prep. 93/94-31, hep-lat 9410011, to appear in Nucl. Phys. B.